

AMISH HOME REMEDIES AND THE RESEARCH SURROUNDING THEM — PART 1

Joshua Kopp, DO

*Primary Investigator, A.T. Still College of Osteopathic Medicine
Psychiatry Resident, Psychiatry Department, Walter Reed National Medical Center*

Lisa Fritz, MD

Research Advisor, A.T. Still College of Osteopathic Medicine



Kopp

This two-part report examines the composition and efficacy of “folk” remedies used by the Amish of Lancaster County. While not a complete list of the plants used, the report seeks to educate medical practitioners regarding Amish practices to help guide patient interviews and prescription recommendations.

INTRODUCTION

An Amish patient with a history of cirrhosis presents to the emergency department with fatigue, shortness of breath, and weight gain. The patient is admitted to the ICU for sepsis complicated by cirrhosis. However, he continues to decline; the patient’s wife believes this is because of antibiotics and gives the patient an extract used by members of the community to aid in liver health.

Soon, the extract bottle is examined by the team, who notes the tincture contains 90% alcohol and may be implicated in the patient’s decline. Once the extract is discontinued, the patient’s liver function tests begin to trend downward, and he can be safely discharged to home health.

This anecdotal story is all too common in Lancaster, Pennsylvania. It illuminates the need to better understand what supplements and alternative treatments patients are taking. The U.S. supplement market generates \$30 billion in yearly revenue, contributing to our community’s health. Exploring alternative therapies may provide opportunities for research.¹

With the ever-changing landscape of modern medicine, it is our duty as practitioners to understand the effects alternative therapies and underutilized treatment options have on our patients. This

review is meant to promote understanding of the home remedies of the Lancaster County Amish community.

METHOD

In this qualitative research, 22 members of the Amish community of Lancaster County were interviewed and ingredients of “natural remedies” were examined to better illuminate home medical practices, especially the composition of some remedies. The interviews took place from February 2022 to July 2024. The individuals interviewed were not asked to provide their names and were assigned an arbitrary number at the beginning of the interview to ensure anonymity.

The interviews were conducted at individuals’ farms, and interviewers began by asking what common ailments were experienced by family members and how ailments are treated at home. Additionally, if known, the active ingredients were cataloged; the literature was subsequently consulted to establish what the scientific community has determined to be the supported indications, contraindications, and adverse effects to consider.

Families were not included if they did not have their home church located in Lancaster County. This was done to ensure consistent information was gathered, as the uses and ingredients of “home remedies” reported by the Lancaster Amish community might vary from those remedies used by others in other locations. This review is in no way comprehensive. Attempts have been made to describe the pertinent ingredients, along with adverse effects, method of action (if known), contraindications, and indications for use.

This study was found to be exempt by the ATSU-Kirksville IRB according to 45CFR46.104 (d)(2)(i). IRB Number: #JK20220314-001.

DISCUSSION

Yarrow Root

Among those interviewed, yarrow root (*Achillea millefolium*, see Fig. 1) was one of the most frequently mentioned plants, noted in 16 of 20 interviews. The genus name is derived from Homer's mention that Achilles used this plant among his troops to stop bleeding.



Fig. 1. Yarrow root (*Achillea millefolium*).

Yarrow contains a pyrrolizidine alkaloid called *achilleine*, which acts as a hemostatic.² It is used by patients in Amish communities for pain, bleeding, and digestion. While clinical studies are sparse, it may decrease perineal pain, edema, and ecchymosis in women undergoing an episiotomy.³

The Amish community uses an ointment containing this extract applied topically. The plant contains caffeoylquinic acid derivatives, which are phytochemicals responsible for antioxidative actions, as well as other chemicals with antibacterial, antiplasmodial, anti-inflammatory, and pain-relieving properties.⁴

It should be noted that yarrow contains the ketone and monoterpene thujone, which can act as a GABA receptor antagonist. This substance has been known to cause a mild, short-lived dermatitis, but is perhaps best known as the component of absinthe once thought to be responsible for causing psychoactive effects; this attribute is no longer thought to be accurate.

When yarrow is used topically, the levels of these substances are not thought to be high enough to cause toxicity.⁴ While more human studies are needed, patients should be educated about the possibility of dermatitis and advised against oral ingestion.

Complete Tissue

Topical creams and balms containing herbals and plant derivatives may be employed by Lancaster County community members to treat infections, back pain, and other ailments. One of these is “complete tissue” or “complete bone and tissue,” which contains white oak bark, marshmallow, black walnut, grave root, skullcap, slippery elm bark, aloe vera, lobelia, vitamin E, beeswax, and wormwood. Black walnut and wormwood are discussed in detail below.

Black Walnut

Black walnut (*Juglans nigra*, see Fig. 2) may also be used for several different problems, including back pain and inflammation, as well as promoting wound healing.

Included in most topical compounds that were identified during this study, black walnut contains many chemicals that appear to be anti-inflammatory and may act as antitumor, antiviral, antioxidant, anti-allergic, antiatherogenic, and antinociceptive; it is thought to reduce cholesterol absorption and work as an antimicrobial against gram-positive bacteria.⁵



Fig. 2. Black walnut (*Juglans nigra*).

assistance would be sought from a licensed physician.

A classic inducer of cytochrome P450 (CYP) enzymes — mainly 1A2, 2D6, and 3A4 — this agent is superior to placebo for the treatment of depression in some trials; it may have a smaller side effect profile than selective serotonin reuptake inhibitors (SSRIs).¹²

Despite some studies showing its efficacy, a systematic review acknowledges that we do not have a full understanding of St. John's wort's side effect profile.¹³ Additionally, although it may be used more frequently in European countries, we do not yet know the optimal daily dosing regimen; long-term studies are lacking.



Fig. 4. St. John's wort (*Hypericum perforatum*).

As noted above, St. John's wort induces CYP enzymes, which can alter the metabolism of drugs like warfarin or digoxin; opiates may also be converted into a more potent form when CYP enzymes are induced. Yet, St. John's wort is well tolerated. A study in 1994 involving 3,250 patients found that only 2.4% had side effects, of which gastrointestinal issues, fatigue, restlessness, and allergic reactions were the most common.¹⁴ St. John's wort is not safe in pregnancy or during lactation.¹⁵

Burdock

Burdock (*Arctium*, see Fig. 5) has been used since ancient times for its potent antioxidant effects. The Amish commonly pick this plant in the spring and dry it to have on hand throughout the year.

It is used topically for wounds, unless there is an accompanying infection, or it can be taken as a tea for gastrointestinal issues, circulation issues, as a diuretic, and for generalized body pain. Although said to be well tolerated, burdock may cause dermatitis, so it is appropriate to watch for erythema.



Fig. 5. Burdock (*Arctium*).

Burdock has many active ingredients, the majority of which act as free radical scavengers.¹⁶ Research suggests it may promote blood circulation to the skin and act as an antidiabetic. It may also act as an inhibitor of pancreatic carcinoma; however, none of these effects have been well studied in humans.¹⁷

Aqueous burdock may treat osteoarthritis by increasing the cellular proliferation of human mesenchymal stem cells, which can, in turn, induce chondrogenesis.¹⁸ Generally, this plant is considered safe, making the risk of misuse and adverse effects extremely low.¹⁷

Horsetail

Horsetail (*Equisetum arvense*, see Fig. 6) is collected and taken for both pain and generalized inflammation. It is a part of many compounds for aches and pains commonly used by members of the Amish community.

This plant has an extremely potent anti-inflammatory action, inhibiting T-cell proliferation and decreasing CD69 and IL-2 surface receptor expression, which in turn decreases production of interferon gamma and tumor necrosis factor alpha.¹⁹

Horsetail should be used cautiously, as it may cause profound hyponatremia.²⁰ Lab work may be indicated if a patient is using horsetail continuously and has abnormal signs and symptoms.

Additionally, case reports implicate horsetail in transient liver function test elevation; horsetail should be avoided in patients with preexisting liver injury or cirrhosis.

CONCLUSION

Often, patients will pursue alternative therapies. To serve communities such as the Amish, drug interactions and proper dosing should be established, with the goal of preventing harm and giving accurate recommendations regarding risks and benefits.

Practitioners should understand these therapies and be prepared to offer recommendations based on the best available evidence. In part 2, we will continue to discuss the compounds introduced in these interviews.




Fig. 6. Horsetail (*Equisetum arvense*).

REFERENCES

- Trinkle KM. Amish culture and their utilization of burns and wounds ointment for the treatment of burns. *Holist Nurs Pract.* 2016;30(2):78-87.
- Macel M. Attract and deter: a dual role for pyrrolizidine alkaloids in plant-insect interactions. *Phytochem Rev.* 2011;10(1):75-82.
- Hajhashemi M, Ghanbari Z, Movahedi M, Rafieian M, Keivani A, Haghollahi F. The effect of *Achillea millefolium* and *Hypericum perforatum* ointments on episiotomy wound healing in primiparous women. *J Matern Fetal Neonatal Med.* 2018;31(1):63-69.
- Czech K, Gawel-Bęben K, Szopa A, et al. Phytochemical profiling, antioxidant and tyrosinase regulatory activities of extracts from herb, leaf and in vitro culture of *Achillea millefolium* (yarrow). *Molecules.* 2023;28(12):4791.
- Ho KV, Schreiber KL, Vu DC, et al. Black walnut (*Juglans nigra*) extracts inhibit proinflammatory cytokine production from lipopolysaccharide-stimulated human promonocytic cell line U-937. *Front Pharmacol.* 2019;10:1059.
- Zhang YZ, Du WX, Fan Y, et al. Purification and characterization of a black walnut (*Juglans nigra*) allergen, Jug n 4. *J Agric Food Chem.* 2017;65(2):454-462.
- Lachenmeier DW. Wormwood (*Artemisia absinthium* L.) – a curious plant with both neurotoxic and neuroprotective properties? *J Ethnopharmacol.* 2010;131(1):224-227.
- Szopa A, Pajor J, Klin P, et al. *Artemisia absinthium* L. – importance in the history of medicine, the latest advances in phytochemistry and therapeutical, cosmetological and culinary uses. *Plants (Basel).* 2020;9(9):1063.
- Ekiert H, Knut E, Świątkowska J, et al. *Artemisia abrotanum* L. (southern Wormwood) – history, current knowledge on the chemistry, biological activity, traditional use and possible new pharmaceutical and cosmetological applications. *Molecules.* 2021;26(9):2503.
- Eadie MJ. Absinthe, epileptic seizures and Valentin Magnan. *J R Coll Physicians Edinb.* 2009;39(1):73-78.
- Mihajilov-Krstev T, Jovanović B, Jović J, et al. Antimicrobial, antioxidative, and insect repellent effects of *Artemisia absinthium* essential oil. *Planta Med.* 2014;80(18):1698-1705.
- Linde K, Berner MM, Kriston L. St. John's wort for major depression. *Cochrane Database Syst Rev.* 2008;2008(4):CD000448.
- Apaydin EA, Maher AR, Shanman R, et al. A systematic review of St. John's wort for major depressive disorder. *Syst Rev.* 2016;5(1):148.
- Woelk H, Burkard G, Grünwald J. Benefits and risks of the hypericum extract LI 160: drug monitoring study with 3250 patients. *J Geriatr Psychiatry Neurol.* 1994;7 Suppl 1:S34-38.
- Avila C, Whitten D, Evans S. The safety of St. John's wort (*Hypericum perforatum*) in pregnancy and lactation: a systematic review of rodent studies. *Phytother Res.* 2018;32(8):1488-1500.
- Parham S, Kharazi AZ, Bakhsheshi-Rad HR, et al. Antioxidant, antimicrobial and antiviral properties of herbal materials. *Antioxidants (Basel).* 2020;9(12):1309.
- Chan YS, Cheng LN, Wu JH, et al. A review of the pharmacological effects of *Arctium lappa* (burdock). *Inflammopharmacol.* 2011;19(5):245-254.
- Wu KC, Weng HK, Hsu YS, Huang PJ, Wang YK. Aqueous extract of *Arctium lappa* L. root (burdock) enhances chondrogenesis in human bone marrow-derived mesenchymal stem cells. *BMC Complement Med Ther.* 2020;20(1):364.
- Gründemann C, Lengen K, Sauer B, Garcia-Käufer M, Zehl M, Huber R. *Equisetum arvense* (common horsetail) modulates the function of inflammatory immunocompetent cells. *BMC Complement Altern Med.* 2014;14:283.
- Norris MR. Severe hyponatremia after drinking horsetail juice. *Ann Intern Med.* 2019;170(9):667-668.

Corresponding Author

Joshua Kopp, DO

Walter Reed National Medical Center

4494 Palmer Rd., Bethesda, MD 20889

(717) 606-6218 / Joshua.W.Kopp@colorado.edu